

## Proximal Hamstring Tendon Repair

### Weeks 0-6:

#### *Goal:*

- Protection of surgical repair

#### *Precautions:*

- Non-weight bearing with crutches for 6 weeks
- No active hamstring contraction
- No hip flexion greater than 45 degrees
- Knee extension limited pending intra-operative tension on the repair

#### *Durable Medical Equipment:*

- Progress ROM by 30 degrees per week to full ROM by 8 weeks
- Cryotherapy unit to be used 4-6 times per day for 20 minutes per session
- Crutches/Non-weight bearing for 4 weeks with progression to full-weight bearing weeks 5-7

#### *Exercises:*

1. Cryotherapy for pain and swelling control 3-5x/day

Teach patient how to transfer from supine to sit, stand to sit, as well as perform ADLs safely. (Avoid greater than 45 degrees of hip flexion when the knee is extended during any and all activities, ie. sitting) Heel props with quad sets, supine position only, to avoid knee stiffness and quad shut down

#### *Other:*

- Light desensitization massage to the incision and posterior hip
- Scar massage
- Silicon patch over incision (if open repair)

#### *Clinical follow-up:*

- Follow-up with physician at approximately 2 weeks postoperative

### Weeks 6-8:

#### *Goal:*

2. Restore normal gait
3. Pain free and normal functional ADLs

#### *Precautions:*

4. Monitor tenderness of surgery site
5. No hamstring flexibility or stretching exercises are to be performed during this phase. Lengthening of the repair and return of normal hamstring flexibility will be allowed to occur on its own. (This is traditionally not an issue following this procedure)

#### *Exercises:*

1. Restore normal gait pattern (emphasize good leg control with extension of knee during swing phase and heel strike)
2. Improve ADL function, ie. Sit -> stand, stairs, etc.

3. Begin light hamstring strengthening with low loads, high reps and high frequency by performing hamstring leg curls in standing with the hip extended. Start with zero resistance then progress as tolerated 1 lb at a time-2sets/20, 4-5x/day
4. Begin total leg strengthening (TLS):
  - a. Heel raises
  - b. Quad sets (active heel lift)
  - c. Short arc quads
  - d. General hip strengthening in side lying (gluteus maximus and medius)
  - e. Single leg balance for proprioception

***Other:***

- Light desensitization massage to the incision and posterior hip
- Scar massage

***Clinical follow-up:***

Follow-up with physician at approximately 8 weeks postoperative to advance rehab and monitor progress

**Weeks 8-12:**

***Goal:***

- Pain-free performance of non-impact aerobic activities
- Unrestricted ADLs at home or work

***Precautions:***

- Monitor hamstring flexibility and tenderness of surgery site

***Exercises:***

1. Begin non-impact aerobic conditioning as tolerated with any of the following:
  - a. Stationary bike
  - b. Stairmaster
  - c. Elliptical trainer
  - d. NordicTrack
  - e. Aquatic therapy with swimming or functional activities in the water (Avoid forceful, explosive or repetitively strainful activities)
2. Continue to progress TLS as tolerated:
  - a. ¼ squats
  - b. Step downs
  - c. Leg press
  - d. Knee extensions
  - e. Heel raises
  - f. Hip abductor in standing with tubing or machine
  - g. Balance and proprioceptive training
3. Progress hamstring strengthening in standing by increasing weight or initiating TheraBand
4. Patient may progress to prone positioning on a machine and then to seated leg curls (with hip flexed at 90 degrees) on a machine or with tubing

***Clinical follow-up:***

Follow-up with physician at approximately 14 weeks postoperative to release to all activities as tolerated. The patient's frequency of follow-up with the therapist will be determined on an individual basis and depend largely on the patient's adherence, as well as the patient's and therapist's comfort level.